



# Cambridge IGCSE™

CANDIDATE NAME



CENTRE NUMBER

--	--	--	--	--

CANDIDATE NUMBER

--	--	--	--

## CHEMISTRY

0620/53

Paper 5 Practical Test

May/June 2025

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions  
Insert (enclosed)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

### INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [ ].
- Notes for use in qualitative analysis are provided in the question paper.

For Examiner's Use	
1	
2	
3	
<b>Total</b>	

This document has **12** pages. Any blank pages are indicated.

\* 00080000002 \*



2

BLANK PAGE



DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN



- 1 You are going to investigate how the rate of reaction between dilute hydrochloric acid and aqueous sodium thiosulfate changes as the concentration of the aqueous sodium thiosulfate decreases. During the reaction, the solution slowly becomes cloudy. As the solution becomes cloudy, it becomes more difficult to see through the solution.


**Read all of the instructions carefully before starting the experiments.**

### Instructions

You are going to do five experiments.

#### Experiment 1

- Use the 50 cm<sup>3</sup> measuring cylinder to pour 50 cm<sup>3</sup> of aqueous sodium thiosulfate into a 100 cm<sup>3</sup> beaker.
- Use the 10 cm<sup>3</sup> measuring cylinder to pour 5 cm<sup>3</sup> of dilute hydrochloric acid into the beaker containing the aqueous sodium thiosulfate.
- Immediately start the stop-watch and stir the contents of the beaker.
- Stand the beaker on the printed insert and look down from above the beaker as shown in Fig. 1.1.



**Fig. 1.1**

- When the text on the printed insert is **not** visible, stop the stop-watch and record the time to the nearest whole number of seconds in Table 1.1.
- Empty the contents of the beaker into the stop-bath and rinse the beaker with distilled water.

#### Experiment 2

- Repeat Experiment 1 using 40 cm<sup>3</sup> of aqueous sodium thiosulfate instead of 50 cm<sup>3</sup>. Add 10 cm<sup>3</sup> of distilled water to the beaker, using the 50 cm<sup>3</sup> measuring cylinder, **before** adding the dilute hydrochloric acid.

#### Experiment 3

- Repeat Experiment 2 using 30 cm<sup>3</sup> of aqueous sodium thiosulfate and 20 cm<sup>3</sup> of distilled water.

#### Experiment 4

- Repeat Experiment 2 using 25 cm<sup>3</sup> of aqueous sodium thiosulfate and 25 cm<sup>3</sup> of distilled water.

#### Experiment 5

- Repeat Experiment 2 using 20 cm<sup>3</sup> of aqueous sodium thiosulfate and 30 cm<sup>3</sup> of distilled water.





(a) Complete Table 1.1.

Table 1.1

experiment	volume of aqueous sodium thiosulfate /cm <sup>3</sup>	volume of distilled water /cm <sup>3</sup>	volume of dilute hydrochloric acid /cm <sup>3</sup>	time taken for text to <b>not</b> be visible/s
1	50	0	5	
2				
3				
4				
5				

[4]

(b) Write a suitable scale on the y-axis and plot your results from Experiments 1 to 5 on Fig. 1.2.

Draw a curve of best fit.




Fig. 1.2

[4]





(c) The relative rate of reaction in each experiment is calculated using the equation shown.

$$\text{relative rate of reaction} = \frac{1}{\text{time taken}}$$

- (i) Calculate the relative rate of reaction in Experiment 1.  
Do **not** give units for your answer.

relative rate of reaction = ..... [1]

- (ii) State in which Experiment, 1, 2, 3, 4 or 5, the relative rate of reaction was greatest.

..... [1]

(d) In each experiment the total volume of aqueous sodium thiosulfate and distilled water is constant.

- (i) Calculate the volume of distilled water needed when the volume of aqueous sodium thiosulfate is 37 cm<sup>3</sup>.

volume = ..... [2]

- (ii) From your graph in Fig. 1.2, deduce the time for the text to **not** be visible when the volume of aqueous sodium thiosulfate is 37 cm<sup>3</sup>.

Show clearly on Fig. 1.2 how you worked out your answer.

time = ..... s [2]

- (iii) Explain why the total volume of aqueous sodium thiosulfate and distilled water is kept constant.

.....

.....

..... [1]

- (e) (i) State why measuring the volume of the dilute hydrochloric acid with a burette instead of a measuring cylinder is an improvement.

.....

..... [1]

- (ii) Explain why it is **not** possible to use a volumetric pipette to measure the volume of aqueous sodium thiosulfate in each experiment.

.....

..... [1]



DO NOT WRITE IN THIS MARGIN



(f) Explain why it is important **not** to change the size of the beaker to a larger beaker during the investigation.

.....  
.....  
..... [2]

[Total: 19]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





2 You are provided with two solids: solid **L** and solid **M**.

Do the following tests on solid **L** and solid **M**. Record all of your observations at each stage.

**Tests on solid L**

Add about 3 cm depth of distilled water to the boiling tube containing solid **L**. Replace the stopper in the boiling tube and shake the boiling tube to dissolve solid **L** and form solution **L**. Divide solution **L** into four approximately equal portions in three boiling tubes and one test-tube.

(a) To the first portion of solution **L** in a boiling tube, add about 2 cm depth of aqueous sodium carbonate.

Record your observations.

.....  
.....  
..... [2]

(b) To the second portion of solution **L** in a boiling tube, add aqueous sodium hydroxide dropwise and then in excess.

Record your observations.

dropwise .....  
in excess ..... [2]

(c) To the third portion of solution **L** in a boiling tube, add aqueous ammonia dropwise and then in excess.

Record your observations.

dropwise .....  
in excess ..... [2]

(d) To the fourth portion of solution **L** in the test-tube, add about 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate.

Record your observations.

.....  
..... [1]

(e) Identify solid **L**.

.....  
..... [2]



DO NOT WRITE IN THIS MARGIN



**Tests on solid M**

**(f)** Carry out a flame test on solid **M**.

Record your observations.

..... [1]

**(g)** Place the remaining solid **M** in a boiling tube. Add about 5 cm depth of distilled water to the boiling tube. Put a stopper in the boiling tube and shake the boiling tube to dissolve solid **M** and form solution **M**. Divide solution **M** into two approximately equal portions in two boiling tubes.

**(i)** To the first portion of solution **M**, add about 1 cm depth of aqueous sodium hydroxide and a piece of aluminium foil. Warm the mixture and test any gas given off.

Record your observations.

.....  
..... [2]

**(ii)** To the second portion of solution **M**, add a few drops of dilute sulfuric acid.

Record your observations.

.....  
..... [1]

**(h)** Identify solid **M**.

.....  
..... [2]

[Total: 15]

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN







\* 00080000010 \*



10

BLANK PAGE

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





## Notes for use in qualitative analysis

### Tests for anions

anion	test	test result
carbonate, $\text{CO}_3^{2-}$	add dilute acid, then test for carbon dioxide gas	effervescence, carbon dioxide produced
chloride, $\text{Cl}^-$ [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
bromide, $\text{Br}^-$ [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	cream ppt.
iodide, $\text{I}^-$ [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate, $\text{NO}_3^-$ [in solution]	add aqueous sodium hydroxide, then aluminium foil; warm carefully	ammonia produced
sulfate, $\text{SO}_4^{2-}$ [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt.
sulfite, $\text{SO}_3^{2-}$	add a small volume of acidified aqueous potassium manganate(VII)	the acidified aqueous potassium manganate(VII) changes colour from purple to colourless

### Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium, $\text{Al}^{3+}$	white ppt., soluble in excess, giving a colourless solution	white ppt., insoluble in excess
ammonium, $\text{NH}_4^+$	ammonia produced on warming	–
calcium, $\text{Ca}^{2+}$	white ppt., insoluble in excess	no ppt. or very slight white ppt.
chromium(III), $\text{Cr}^{3+}$	green ppt., soluble in excess	green ppt., insoluble in excess
copper(II), $\text{Cu}^{2+}$	light blue ppt., insoluble in excess	light blue ppt., soluble in excess, giving a dark blue solution
iron(II), $\text{Fe}^{2+}$	green ppt., insoluble in excess, ppt. turns brown near surface on standing	green ppt., insoluble in excess, ppt. turns brown near surface on standing
iron(III), $\text{Fe}^{3+}$	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc, $\text{Zn}^{2+}$	white ppt., soluble in excess, giving a colourless solution	white ppt., soluble in excess, giving a colourless solution





### Tests for gases

gas	test and test result
ammonia, $\text{NH}_3$	turns damp red litmus paper blue
carbon dioxide, $\text{CO}_2$	turns limewater milky
chlorine, $\text{Cl}_2$	bleaches damp litmus paper
hydrogen, $\text{H}_2$	'pops' with a lighted splint
oxygen, $\text{O}_2$	relights a glowing splint
sulfur dioxide, $\text{SO}_2$	turns acidified aqueous potassium manganate(VII) from purple to colourless

### Flame tests for metal ions

metal ion	flame colour
lithium, $\text{Li}^+$	red
sodium, $\text{Na}^+$	yellow
potassium, $\text{K}^+$	lilac
calcium, $\text{Ca}^{2+}$	orange-red
barium, $\text{Ba}^{2+}$	light green
copper(II), $\text{Cu}^{2+}$	blue-green

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

