



# Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**FURTHER MATHEMATICS**

**9231/33**

Paper 3 Further Mechanics

**October/November 2020**

**1 hour 30 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

## INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ ms}^{-2}$ .

## INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Blank pages are indicated.



- 1 A particle  $P$  of mass  $m$  is placed on a fixed smooth plane which is inclined at an angle  $\theta$  to the horizontal. A light spring, of natural length  $a$  and modulus of elasticity  $3mg$ , has one end attached to  $P$  and the other end attached to a fixed point  $O$  at the top of the plane. The spring lies along a line of greatest slope of the plane. The system is released from rest with the spring at its natural length.

Find, in terms of  $a$  and  $\theta$ , an expression for the greatest extension of the spring in the subsequent motion. [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



- 3 One end of a light elastic string, of natural length  $a$  and modulus of elasticity  $4mg$ , is attached to a fixed point  $O$ . The other end of the string is attached to a particle of mass  $m$ . The particle moves in a horizontal circle with a constant angular speed  $\sqrt{\frac{g}{a}}$  with the string inclined at an angle  $\theta$  to the downward vertical through  $O$ . The length of the string during this motion is  $(k+1)a$ .

- (a) Find the value of  $k$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the value of  $\cos \theta$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....



The diagram shows the cross-section  $ABCD$  of a uniform solid object which is formed by removing a cone with cross-section  $DCE$  from the top of a larger cone with cross-section  $ABE$ . The perpendicular distance between  $AB$  and  $DC$  is  $h$ , the diameter  $AB$  is  $6r$  and the diameter  $DC$  is  $2r$ .

- (a) Find an expression, in terms of  $h$ , for the distance of the centre of mass of the solid object from  $AB$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- 5 A particle  $P$  is projected with speed  $u$  at an angle  $\alpha$  above the horizontal from a point  $O$  on a horizontal plane and moves freely under gravity. The horizontal and vertical displacements of  $P$  from  $O$  at a subsequent time  $t$  are denoted by  $x$  and  $y$  respectively.

- (a) Derive the equation of the trajectory of  $P$  in the form

$$y = x \tan \alpha - \frac{gx^2}{2u^2} \sec^2 \alpha. \quad [3]$$

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

The point  $Q$  is the highest point on the trajectory of  $P$  in the case where  $\alpha = 45^\circ$ .

- (b) Show that the  $x$ -coordinate of  $Q$  is  $\frac{u^2}{2g}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....







A series of horizontal dotted lines for writing, spanning the width of the page.









**BLANK PAGE**

---

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at [www.cambridgeinternational.org](http://www.cambridgeinternational.org) after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.