



## Cambridge International AS & A Level

CANDIDATE  
NAME

--

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**MATHEMATICS**

**9709/42**

Paper 4 Mechanics

**October/November 2022**

**1 hour 15 minutes**

You must answer on the question paper.

You will need: List of formulae (MF19)

### INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity ( $g$ ) is needed, use  $10 \text{ m s}^{-2}$ .

### INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **12** pages.



2 A particle  $P$  of mass  $0.4\text{ kg}$  is in limiting equilibrium on a plane inclined at  $30^\circ$  to the horizontal.

(a) Show that the coefficient of friction between the particle and the plane is  $\frac{1}{3}\sqrt{3}$ . [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A force of magnitude  $7.2\text{ N}$  is now applied to  $P$  directly up a line of greatest slope of the plane.

(b) Given that  $P$  starts from rest, find the time that it takes for  $P$  to move  $1\text{ m}$  up the plane. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

3



A particle of mass 0.3 kg is held at rest by two light inextensible strings. One string is attached at an angle of  $60^\circ$  to a horizontal ceiling. The other string is attached at an angle  $\alpha^\circ$  to a vertical wall (see diagram). The tension in the string attached to the ceiling is 4 N.

Find the tension in the string which is attached to the wall and find the value of  $\alpha$ . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

4 A car of mass 1200 kg is travelling along a straight horizontal road  $AB$ . There is a constant resistance force of magnitude 500 N. When the car passes point  $A$ , it has a speed of  $15 \text{ m s}^{-1}$  and an acceleration of  $0.8 \text{ m s}^{-2}$ .


(a) Find the power of the car's engine at the point  $A$ . [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

The car continues to work with this power as it travels from  $A$  to  $B$ . The car takes 53 seconds to travel from  $A$  to  $B$  and the speed of the car at  $B$  is  $32 \text{ m s}^{-1}$ .

(b) Show that the distance  $AB$  is 1362.6 m. [3]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....



A block A of mass 80 kg is connected by a light, inextensible rope to a block B of mass 40 kg. The rope joining the two blocks is taut and is parallel to a line of greatest slope of a plane which is inclined at an angle of  $20^\circ$  to the horizontal. A force of magnitude 500 N inclined at an angle of  $15^\circ$  above the same line of greatest slope acts on A (see diagram). The blocks move up the plane and there is a resistance force of 50 N on B, but no resistance force on A.

- (a) Find the acceleration of the blocks and the tension in the rope. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

**(b)** Find the time that it takes for the blocks to reach a speed of  $1.2 \text{ m s}^{-1}$  from rest. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 6 Three particles  $A$ ,  $B$  and  $C$  of masses  $0.3\text{ kg}$ ,  $0.4\text{ kg}$  and  $m\text{ kg}$  respectively lie at rest in a straight line on a smooth horizontal plane. The distance between  $B$  and  $C$  is  $2.1\text{ m}$ .  $A$  is projected directly towards  $B$  with speed  $2\text{ m s}^{-1}$ . After  $A$  collides with  $B$  the speed of  $A$  is reduced to  $0.6\text{ m s}^{-1}$ , still moving in the same direction.

- (a) Show that the speed of  $B$  after the collision is  $1.05\text{ m s}^{-1}$ . [2]

.....

.....

.....

.....

.....

.....

.....

After the collision between  $A$  and  $B$ ,  $B$  moves directly towards  $C$ . Particle  $B$  now collides with  $C$ . After this collision, the two particles coalesce and have a combined speed of  $0.5\text{ m s}^{-1}$ .

- (b) Find  $m$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



- (c) Find the time that it takes, from the instant when  $B$  and  $C$  collide, until  $A$  collides with the combined particle. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

7 A particle  $P$  travels in a straight line, starting at rest from a point  $O$ . The acceleration of  $P$  at time  $t$  s after leaving  $O$  is denoted by  $a \text{ m s}^{-2}$ , where

$$a = 0.3t^{\frac{1}{2}} \quad \text{for } 0 \leq t \leq 4,$$
$$a = -kt^{-\frac{3}{2}} \quad \text{for } 4 < t \leq T,$$

where  $k$  and  $T$  are constants.

(a) Find the velocity of  $P$  at  $t = 4$ . [2]

.....

.....

.....

.....

.....

.....

.....

.....

(b) It is given that there is no change in the velocity of  $P$  at  $t = 4$  and that the velocity of  $P$  at  $t = 16$  is  $0.3 \text{ m s}^{-1}$ .

Show that  $k = 2.6$  and find an expression, in terms of  $t$ , for the velocity of  $P$  for  $4 \leq t \leq T$ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Given that  $P$  comes to instantaneous rest at  $t = T$ , find the exact value of  $T$ . [2]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

(d) Find the total distance travelled between  $t = 0$  and  $t = T$ . [4]

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

